

Orientable smooth manifolds are essentially quasigroups

Charlotte Aten (joint work with Semin Yoo)

University of Rochester

2021 November 7

Introduction

- In the mid-2010s Herman and Pakianathan introduced a functorial construction of closed surfaces from noncommutative finite groups.
- Semin Yoo and I decided to produce an n -dimensional generalization.
- The two main challenges in doing this were finding an appropriate analogue of noncommutative groups and in desingularizing the n -dimensional pseudomanifolds which arose at the first stage of our construction.
- Ultimately we found that every orientable triangulable manifold could be manufactured in the manner we described.

Talk outline

- Herman and Pakianathan's construction
- Quasigroups
- The first functor: Open serentation
- The second functor: Serentation

Herman and Pakianathan's construction

- Consider the quaternion group \mathbf{G} of order 8 whose universe is $G := \{\pm 1, \pm i, \pm j, \pm k\}$.
- We begin by picking out all the pairs of elements $(x, y) \in G^2$ so that $xy \neq yx$. We call this collection $\text{NCT}(\mathbf{G})$.
- We define $\text{In}(\mathbf{G})$ to be all the elements of G which are entries in some pair $(x, y) \in \text{NCT}(\mathbf{G})$.
- Similarly, $\text{Out}(\mathbf{G})$ is defined to be all the members of G of the form $f(x, y)$ where $(x, y) \in \text{NCT}(\mathbf{G})$.

Herman and Pakianathan's construction

- In this case we have

$$\text{NCT}(\mathbf{G}) = \left\{ (\pm u, \pm v) \mid \{u, v\} \in \binom{\{i, j, k\}}{2} \right\}$$

so

$$\text{In}(\mathbf{G}) = \{\pm i, \pm j, \pm k\}$$


and

$$\text{Out}(\mathbf{G}) = \{\pm i, \pm j, \pm k\}.$$

- From this data we form a simplicial complex (actually a 2-pseudomanifold) whose facets are of the form $\{\underline{x}, \underline{y}, \overline{f(x, y)}\}$ where $(x, y) \in \text{NCT}(\mathbf{G})$.

Herman and Pakianathan's construction

- During the talk I drew a part of this complex here:




Herman and Pakianathan's construction

- The three 4-cycles

$$(\underline{i}, \underline{j}, \underline{-i}, \underline{-j}), (\underline{i}, \underline{k}, \underline{-i}, \underline{-k}), \text{ and } (\underline{j}, \underline{k}, \underline{-j}, \underline{-k}).$$

each carry an octahedron.



Herman and Pakianathan's construction

- This simplicial complex, which we call **Sim(G)** and Herman and Pakianathan called $X(Q_8)$, consists of three 2-spheres, each pair of which is glued at two points.
- Deleting these points to disjointize the spheres and filling the resulting holes yields the manifold we call **Ser(G)** and Herman and Pakianathan called $Y(Q_8)$.
- In this case **Ser(G)** is the disjoint union of three 2-spheres.

Quasigroups

Definition (Quasigroup)

A (*binary*) *quasigroup* is a magma $\mathbf{A} := (A, f: A^2 \rightarrow A)$ such that if any two of the variables x , y , and z are fixed the equation

$$f(x, y) = z$$

has a unique solution.

- That is, a quasigroup is a magma whose Cayley table is a Latin square, where each entry occurs once in each row and each column.
- All groups are quasigroups, but quasigroups need not have identities or be associative.

Quasigroups

- The midpoint operation

$$f(x, y) := \frac{1}{2}(x + y)$$

is a quasigroup operation on \mathbb{R}^n .

- The magma $(\mathbb{Z}, -)$ is a quasigroup.

Quasigroups

Definition (Quasigroup)

A (binary) quasigroup is an algebra $\mathbf{A} := (A, f, g_1, g_2)$ where for all $x_1, x_2, y \in A$ we have

$$f(g_1(x_2, y), x_2) = y,$$

$$f(x_1, g_2(x_1, y)) = y,$$

$$g_1(x_2, f(x_1, x_2)) = x_1,$$

and

$$g_2(x_1, f(x_1, x_2)) = x_2.$$

- We think of $g_1(x, y)$ as the division of y by x in the second coordinate.

Quasigroups

- The preceding definition shows that the class Quas_2 of all binary quasigroups can be defined by universally-quantified equations, or *identities*.
- This means that Quas_2 is a variety of algebras in the sense of universal algebra, and hence forms a category \mathbf{Quas}_2 which is closed under taking quotients, subalgebras, and products.
- Note that Herman and Pakianathan's construction works with noncommutative quasigroups just as well as with groups.
- We would then like an n -ary version of a quasigroup for our n -dimensional generalization.

Quasigroups

Definition (Quasigroup)

An n -quasigroup is an n -magma $\mathbf{A} := (A, f: A^n \rightarrow A)$ such that if any $n - 1$ of the variables x_1, \dots, x_n, y are fixed the equation

$$f(x_1, \dots, x_n) = y$$

has a unique solution.

- That is, an n -quasigroup is an n -magma whose Cayley table is a Latin n -cube.
- All n -ary groups are quasigroups, but quasigroups need not be associative.

Quasigroups

- Given any group \mathbf{G} the n -ary multiplication

$$f(x_1, \dots, x_n) := x_1 \cdots x_n$$

is a quasigroup operation on G .

Quasigroups

Definition (Quasigroup)

An n -quasigroup is an algebra

$$\mathbf{A} := (A, f, g_1, \dots, g_n)$$

where for all $x_1, \dots, x_n, y \in A$ and each $i \in \{1, 2, \dots, n\}$ we have

$$f(x_1, \dots, x_{i-1}, g_i(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n, y), x_{i+1}, \dots, x_n) = y$$

and

$$g_i(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n, f(x_1, \dots, x_n)) \approx x_i.$$

- We think of $g_i(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n, y)$ as the division of y simultaneously by x_j in the j^{th} coordinate for each $j \neq i$.

Quasigroups

- We say that an n -quasigroup \mathbf{A} is *commutative* when for all $x_1, \dots, x_n \in A$ and all $\sigma \in \text{Perm}_n$ we have

$$f(x_1, \dots, x_n) = f(x_{\sigma(1)}, \dots, x_{\sigma(n)}).$$

- We say that an n -quasigroup \mathbf{A} is *alternating* when for all $x_1, \dots, x_n \in A$ and all $\sigma \in \text{Alt}_n$ we have

$$f(x_1, \dots, x_n) = f(x_{\sigma(1)}, \dots, x_{\sigma(n)}).$$

- Our “correct” analogue of the variety of groups will be the variety AQ_n of alternating n -ary quasigroups.

Quasigroups

- By general results in universal algebra there are nontrivial members of AQ_n for each n , but the easiest examples are either commuting (take the n -ary multiplication for an abelian group) or infinite (the free alternating quasigroups, which we need later but are too much right now).
- We tediously found the following example by hand:

Quasigroups

- Take $S := (\mathbb{Z}/5\mathbb{Z})^3$ and define $h: \mathbb{Z}/5\mathbb{Z} \times \mathbf{Alt}_3 \rightarrow \mathbf{Perm}_S$ by

$$(h(k, \sigma))(x_1, x_2, x_3) := (x_{\sigma(1)} + k, x_{\sigma(2)} + k, x_{\sigma(3)} + k).$$

There are 7 members of $\text{Orb}(h)$. One system of orbit representatives is:

$$\{000, 011, 022, 012, 021, 013, 031\}.$$

Quasigroups

- Let $A := \mathbb{Z}/5\mathbb{Z}$ and define a ternary operation $f: A^3 \rightarrow A$ so that

$$f((h(k, \sigma))(x_1, x_2, x_3)) = f(x_1, x_2, x_3) + k$$

and f is defined on the above set of orbit representatives as follows.

xyz	$f(x, y, z)$
000	0
011	0
022	0
012	3
021	4
013	4
031	2

Quasigroups

- By taking products of $\mathbf{A} := (A, f)$ this gives us infinitely many finite, noncommutative, alternating ternary quasigroups, but we only have one basic example.
- We reached out to Jonathan Smith to see if anyone had studied the varieties of alternating n -quasigroups before, but it seemed that no one had.
- He did, however, give us an example which we generalized into an *alternating product* construction which takes an n -ary commutative quasigroup and an $(n + 1)$ -ary commutative quasigroup and yields an n -ary alternating quasigroup which is typically not commutative.

The first functor: Open serenation

- Our first construction gives a functor

$$\mathbf{OSer}_n: \mathbf{NCAQ}_n \rightarrow \mathbf{SMfld}_n.$$

- We define

$$\mathbf{Sim}_n: \mathbf{NCAQ}_n \rightarrow \mathbf{PMfld}_n$$

similarly to our previous example for $n = 2$.

- We define $\mathbf{NCT}(\mathbf{A})$ to consist of all tuples $(a_1, \dots, a_n) \in A^n$ such that $f(a_1, \dots, a_n) \neq f(a_2, a_1, \dots, a_n)$.
- We define $\mathbf{In}(\mathbf{A})$ to consist of all entries in noncommuting tuples of \mathbf{A} and $\mathbf{Out}(\mathbf{A})$ to consist of all $f(a_1, \dots, a_n)$ where $(a_1, \dots, a_n) \in \mathbf{NCT}(\mathbf{A})$.

The first functor: Open serenation

- We set

$$\text{Sim}(\mathbf{A}) := \{ \underline{a} \mid a \in \text{In}(\mathbf{A}) \} \cup \{ \bar{a} \mid a \in \text{Out}(\mathbf{A}) \}$$

and

$$\text{SimFace}(\mathbf{A}) := \bigcup_{a \in \text{NCT}(\mathbf{A})} \text{Sb} \left(\left\{ \underline{a}_1, \dots, \underline{a}_n, \overline{f(a)} \right\} \right).$$

- We define

$$\mathbf{Sim}_n(\mathbf{A}) := (\text{Sim}(\mathbf{A}), \text{SimFace}(\mathbf{A})).$$

The first functor: Open serenation

- We create $\mathbf{OSer}_n(\mathbf{A})$ by taking the open geometric realization of $\mathbf{Sim}_n(\mathbf{A})$ (basically all but the $(n - 2)$ -skeleton of the open geometric realization) and then equipping it with a smooth atlas.
- The *standard open bipyramid* (or just *bipyramid*) in \mathbb{R}^n is

$$\text{Bipyr}_n := \text{OCvx} \left(\left\{ (0, \dots, 0), \left(\frac{2}{n}, \dots, \frac{2}{n} \right) \right\} \cup \{e_1, \dots, e_n\} \right)$$

where e_i is the i^{th} standard basis vector of \mathbb{R}^n .

The first functor: Open seriation

- Given an alternating n -quasigroup \mathbf{A} and $a = (a_1, \dots, a_n) \in \text{NCT}(\mathbf{A})$ the *serene chart* of input type for a is

$$\underline{\phi}_a : \text{Bipyr}_n \rightarrow \text{OSer}_n(\mathbf{A}).$$

- We set

$$\underline{\phi}_a(u_1, \dots, u_n) := \sum_{i=1}^n u_i \underline{a}_i + \left(1 - \sum_{i=1}^n u_i\right) \overline{f(a)}$$

when $\sum_{i=1}^n u_i \leq 1$.

- Otherwise,

$$\underline{\phi}_a(u_1, \dots, u_n) := \frac{2}{n} \sum_{i=1}^n \left(1 + \frac{n-2}{2} u_i - \sum_{j \neq i} u_j\right) \underline{a}_i + \left(-1 + \sum_{i=1}^n u_i\right) \overline{f(a')}.$$

The first functor: Open serenation

- There are also serene charts of output type, where are defined similarly.
- We set

$$(\mathbf{OSer}_n(\mathbf{A}), \tau) := (\mathbf{OGeo}_n \circ \mathbf{Sim}_n)(\mathbf{A}).$$

- We then define


$$\mathbf{OSer}_n(\mathbf{A}) := (\mathbf{OSer}_n(\mathbf{A}), \tau, \mathbf{SerAt}_n(\mathbf{A}))$$

where

$$\mathbf{SerAt}_n(\mathbf{A}) := \bigcup_{a \in \mathbf{NCT}(\mathbf{A})} \{ \underline{\phi}_a, \overline{\phi}_a \}.$$


The first functor: Open serenation

- The incidence graph of the facets of $\mathbf{Sim}(\mathbf{A})$ for the ternary quasigroup \mathbf{A} from the previous example is pictured.



The first functor: Open serenation

- The 1-skeleton of $\mathbf{Sim}(\mathbf{A})$ for the ternary quasigroup \mathbf{A} from the previous example is pictured.



The first functor: Open serenation

- One may verify that $\mathbf{OSer}(\mathbf{A})$ is a 3-sphere minus the graph pictured previously, which is homotopy equivalent to the join of 21 circles.

The second functor: Serenation

- For any alternating quasigroup \mathbf{A} we may equip $\mathbf{OSer}(\mathbf{A})$ with a Riemannian metric in a functorial manner which makes $\mathbf{OSer}(\mathbf{A})$ flat.
- We then define a *Euclidean metric completion functor*

$$\mathbf{EuCmplt}: \mathbf{Riem}_n \rightarrow \mathbf{Mfld}_n$$

which assigns to a Riemannian manifold (\mathbf{M}, g) the topological manifold consisting of all points in the metric completion of \mathbf{M} which are locally Euclidean.

The second functor: Serenation

- The *serenation functor*

$$\mathbf{Ser}_n: \mathbf{NCAQ}_n \rightarrow \mathbf{Mfld}_n$$

is given by

$$\mathbf{Ser}(\mathbf{A}) := \mathbf{EuCmplt}(\mathbf{OSer}(\mathbf{A}), g)$$

where g is the standard metric on $\mathbf{OSer}(\mathbf{A})$.

- In the previous example of the ternary quasigroup \mathbf{A} we find that $\mathbf{Ser}_3(\mathbf{A})$ is the 3-sphere.

The second functor: Serenation

Definition (Serene manifold)

We say that a connected orientable n -manifold \mathbf{M} is *serene* when there exists some alternating n -quasigroup \mathbf{A} such that \mathbf{M} is a component of $\mathbf{Ser}(\mathbf{A})$.

The second functor: Serenation

Theorem (A., Yoo (2021))

Every connected orientable triangulable n -manifold is serene.

The second functor: Serenation

Theorem (A., Yoo (2021))

Every connected orientable triangulable n -manifold is serene.

- Consider a triangulation of the given manifold \mathbf{M} .
- Subdivide each facet in a manner I will draw off to the side.
- We have that \mathbf{M} is homeomorphic to a corresponding component of the serenation of a quotient of the free alternating n -quasigroup whose generators are the vertices of the subdivided triangulation.

References

- Mark Herman and Jonathan Pakianathan. “On a canonical construction of tessellated surfaces from finite groups”. In: *Topology Appl.* 228 (2017), pp. 158–207. ISSN: 0166-8641